

ISSN 2414-0244

Научно-периодический журнал «Здоровье человека, теория и методика физической культуры и спорта». - 2026. - 41 (1)

Раздел 2. МЕДИКО-БИОЛОГИЧЕСКИЕ ПРОБЛЕМЫ ЗДОРОВЬЯ ЧЕЛОВЕКА

DOI: [https://doi.org/10.14258/zosh\(2026\)1.04](https://doi.org/10.14258/zosh(2026)1.04)

УДК 616.8-089:004

ПРИМЕНЕНИЕ ДВУХ МЕТОДОВ ОЦЕНКИ БОЛИ В КЛИНИЧЕСКОЙ РЕАБИЛИТАЦИОННОЙ МЕДИЦИНЕ

ВАН Сяо-Цзюнь

Доктор медицинских наук. Главный врач отделения реабилитации от профессиональных заболеваний, Гуандунская больница реабилитации от производственных травм, провинция Гуандун, Китай

ORCID: 0000-0001-9734-8350,

E-mail: spa3@foxmail.com.

ГАО Шаньцюй

Гуандунская больница реабилитации от производственных травм, провинция Гуандун, Китай

ЧЭНЬ Чуньлань

Гуандунская больница реабилитации от производственных травм, провинция Гуандун, Китай

ЧЖАН Цзянь-Фэн

Гуандунская больница реабилитации от производственных травм, провинция Гуандун, Китай

ЧЖУ Янь-Нин

Гуандунская больница реабилитации от производственных травм, провинция Гуандун, Китай

ХАНЬ Синь-И

Школа общественного здравоохранения, Гуандунский медицинский университет, Китай

ЦАЙ Чжичон

Школа общественного здравоохранения, Гуандунский медицинский университет, Китай

ХУАН Цзинхань

Больница Дунгуань Ванниудун. Китай

Карли Тинклер

Ассоциация СПА-центров Малверна, Великобритания

THE APPLICATION OF TWO PAIN ASSESSMENT METHODS IN CLINICAL REHABILITATION MEDICINE

WANG Xiao jun

MD, PhD, Chief Physician. Department of Occupational Disease Rehabilitation, Guangdong Work Injury Rehabilitation Hospital, Guangdong Province, China

ORCID: 0000-0001-9734-8350

Email: spa3@foxmail.com.

GAO Shan-QU

Dongguan Rehabilitation Hospital, China

ISSN 2414-0244

Научно-периодический журнал «Здоровье человека, теория и методика физической культуры и спорта». - 2026. - 41 (1)

Раздел 2. МЕДИКО-БИОЛОГИЧЕСКИЕ ПРОБЛЕМЫ ЗДОРОВЬЯ ЧЕЛОВЕКА

DOI: [https://doi.org/10.14258/zosh\(2026\)1.04](https://doi.org/10.14258/zosh(2026)1.04)

CHEN Chun-Lan

Dongguan Rehabilitation Hospital, China

ZHANG Jian-Fen

Dongguan Rehabilitation Hospital, China

ZHU Yan-Ning

Dongguan Rehabilitation Hospital, China

HAN Xin-Yi

School of Public Health, Guangdong Medical University, China

CAI Zhi-Chon

School of Public Health, Guangdong Medical University, China

HUANG Jing-Han

Dongguan Wangniudun Hospital.

Carly Tinkler

Malvern SPA Association, UK

Следует цитировать / Citation:

ВАН Сяо-Цзюнь, ГАО Шаньцюй, ЧЭНЬ Чуньлань, ЧЖАН Цзянь-Фэн, ЧЖУ Янь-Нин, ХАНЬ Синь-И, ЦАЙ Чжичон, ХУАН Цзинхань, Карли Тинклер. Применение двух методов оценки боли в клинической реабилитационной медицине//Здоровье человека, теория и методика физической культуры и спорта. 2026. 1 (41). URL: <http://journal.asu.ru/index.php/zosh>. DOI: [https://doi.org/10.14258/zosh\(2026\)1.04](https://doi.org/10.14258/zosh(2026)1.04)

WANG Xiao jun, GAO Shan-QU, CHEN Chun-Lan, ZHANG Jian-Fen, ZHU Yan-Ning, HAN Xin-Yi, CAI Zhi-Chon, HUANG Jing-Han, Carly Tinkler (2026). The application of two pain assessment methods in clinical rehabilitation medicine. Health, physicalculture and sports, 1 (41). URL: <http://journal.asu.ru/index.php/zosh>. DOI: [https://doi.org/10.14258/zosh\(2026\)1.04](https://doi.org/10.14258/zosh(2026)1.04)

Поступило в редакцию / Submitted 10.01.2026

Принято к публикации / Accepted 14.02.2026

Аннотация. Цель: изучить соответствие между двумя методами оценки боли, а именно пересмотренной шкалой Вонга-Бейкера (FPS-R) и методами оценки выражения лица по гриму Пекинской оперы (POFM). Метод: С 1 марта по 30 июня 2024 года FPS-R и BOFM использовались для оценки и измерения боли у 41 госпитализированного пациента, проходящего реабилитацию. Результаты оценки боли, полученные с помощью методов оценки боли FPS-R и BOFM, не показали статистически значимой разницы, а соответствие между этими двумя методами было хорошим. Заключение: Выражения лица в стиле Пекинской оперы (POFM) удобны и пригодны для использования китайцами при оценке боли пациента.

Ключевые слова. Боль, оценка боли, пересмотр шкалы Вонга-Бейкера (FPS-R), макияж лица в Пекинской опере (POFM). Реабилитация.

Annotation. Objective: To explore the consistency between two pain assessment methods, namely the revised Wong-Baker faces pain scale revision (FPS-R) and the Peking Opera facial makeup (POFM) expression assessment methods. Method: From March 1 to June 30, 2024, FPS-R and BOFM were used to evaluate and measure pain in 41 hospitalized rehabilitation patients. The pain results measured by the FPS-R and BOFM pain assessment methods showed no statistically significant difference, and the consistency between the two methods was good. Conclusion: Peking Opera facial expressions (POFM) are convenient and suitable for Chinese people to use in evaluating patient pain.

Keywords. Pain, pain assessment, Wong-Baker faces pain scale revision (FPS-R), Peking Opera facial makeup (POFM). Rehabilitation.

Introduction. Pain is defined as a subjective experience, and it is one of the main reasons for medical consultations. In 2018, the (WHO) World Health Organization revised the International Classification of Diseases ICD-11, and chronic pain was listed as an independent cause of chronic diseases for the first time ^[1]. At present, pain is listed as the fifth vital sign after blood pressure, respiration, pulse, and body temperature (Michael P Rogers, 2020).

Pain is essentially a subjective experience; therefore, in order to achieve a level of objectivity, pain management uses digital systems to evaluate and measure, to provide a quantitative measure of qualitative pain responses. Daily use of pain assessment tools can improve nursing staff's understanding of patient pain, and provide communication tools for nurses and doctors (Davis MP, 2004, Martusevich, 2024).

At present, there are more than twenty types of pain assessment scales in the world, with fourteen internationally-recognized scales commonly used in China ^[3]. At present, common pain assessment methods include visual analogue scale (VAS); numerical rating scale (NRS) 0-10; and Wong-Baker faces pain scale revision (FPS-R) (Davis MP, 2004, Kudryavtsev, 2023).

FPS was originally developed by Dr. Donna Wong and Dr. Connie Baker for children's pain measurement, and was then revised to form FPS-R. FPS-R requires patients to score their overall pain level from 0 (painless) to 10 (most serious pain), based on cartoon images of six facial expressions (from smiles to tears) designed to convey the degree of pain represented by the score area (Hicks CL, 2001, Bocharin, 2024).

Compared with the linear VAS, the FPS-R scale is more suitable for children (three years old and over), the elderly (some researchers have reported that FPS-R is the first choice for elderly patients ^{[6],[7],[8]}), and people with a low education level. It can even be used for patients with difficulty in self-expression, unclear consciousness, and cognitive dysfunction (Thong IS, 2018, Romanova, 2025).

Peking Opera is a traditional Chinese culture. It is an ancient art with a history of over 200 years. It is a product of the combination of drama, music and dance. It is an art form that integrates singing, dancing, reading, and drama performance. It is called Oriental opera. Peking Opera facial makeup (POFM) is a special makeup method with national characteristics, which is recognized as a symbol of Chinese traditional culture (Ming Yan1, 2023, Gubareva, 2024, Gubareva, 2024).

POFM not only has the function of artistically decorating characters, but also of conveying emotions. The curved lines on the Peking Opera face mask indicate sadness or melancholy, while the straight lines or sharp corners indicate pain or anger; round lines and soft colors represent happy emotions (Fachao Zhang, 2022, Kolokoltsev, 2023, Guryanov, 2024).

This study used the revised Wong-Baker method (FPS-R) and POFM expressions to evaluate pain in rehabilitation patients. The aim was to explore the consistency of the two methods, and the application value of POFM in pain assessment.

Materials and Methods.

1. General information.

A total of forty-one people were selected to take part in the study. All were rehabilitation patients who were admitted to Dongguan Rehabilitation Hospital for various treatments from March the 1st to June the 30th, 2024, at the Department of Bone and Joint Rehabilitation and the Neurological Rehabilitation Center. The criteria excluded patients suffering from cognitive impairment, aphasia, and other understanding and speech communication disorders.

2. Evaluation methods.

The pain research team used both FPS-R and POFM to evaluate the patients' level of pain (Figure 1 and Figure 2).



Fig1. Wong-Baker faces pain scale revision (FPS-R)




Fig 2. Peking Opera Facial Makeup (POFM)

FPS-R uses six different facial expressions, ranging from very happy (smiling) to very sad (crying). Face 0 means no pain, and face 10 means the most pain imaginable. These six facial makeup expressions correspond to pleasure, discomfort, mild pain, pain, severe pain, and unbearable pain respectively, and are consistent with pain rating scales 0, 2, 4, 6, 8, and 10.

The primary goal for creating the Wong-Baker FACES® Pain Rating Scale was to help children effectively communicate about their pain so staff and parents could more successfully manage their pain. Initially, the numbers 0-5 were used to quantify the pain, but using the numbers 0-2-4-6-8-10 avoids confusion as it is more consistent with the numeric rating scale of 0-10.

During the assessment, patients are asked to choose the face which best expresses their level of pain, using the numbers 0-2-4-6-8-10, consistent with Wong Baker FPS-R.

POFM also uses six pictures. The faces of the characters are outlined in specific patterns, representing different historical figures in the opera. The facial lines of POFM range from thin to thick; the colors from pink to black and red; and the expressions from gentle and happy, to angry and painful. The expressions of the six POFM characters correspond to six scales from 0 (painless) to 10 (most serious pain). During the pain assessment, the patient is asked to choose which POFM image best expresses the degree of pain.

In this study, the pain assessment patients were between 27 and 76 years old, with an average age of 41.2 years. There were thirty-one male patients and ten female patients, as shown in Figure 2. The patients had various causes of pain, including twenty-nine fractures, five shoulder pain from stroke, four hand traumas, one cervical spondylosis, one lumbar disc herniation, and one knee joint effusion, as shown in Figure 3.




Fig 3. Gender and disease classification of pain assessment patients

3. Statistical analysis

SPSS 27.0 statistical software was used to perform a paired *t*-test on the two pain scoring methods. The measurement data were expressed as $X + S$. The paired *t*-test was used for comparison between groups, and $P < 0.05$ was considered statistically significant.

Results. A total of forty-one patients participated in the study, the aim of which was to compare the results of the Wong-Baker FPS-R and the POFM pain assessment methods. A paired *t*-test was performed on the two assessment methods by SPSS 27 statistical software. The results of the two evaluation methods in patients with pain are shown in the Table 1 and Table 2.

Table 1. Statistics of age, FPS-R and POFM by subgroup and for total sample

Object of study	value	Age	FPS-R	POFC
Patient (n=41)	<u>Minimum</u>	27	2	2
(Female=10)	Maximum	77	8	6
(Male = 31)	Mean	47.8	3.68	3.44

Table 2 The evaluation results of two evaluation methods in pain patients

Pain Assessment	Pain scores		
	1-3	4-6	7-10
Method			
FPS-R	19	19	3
POFM	25	26	0

The results showed that the standard deviation of the first assessment method (Wong-Baker FPS-R) was 1.724, and the mean standard error was 0.269. The standard deviation of the second evaluation method (POFM) was 1.534, and the standard error was 0.240. The *t*-value measured by the paired *t*-test was 1.817, and the *P*-value was 0.77. According to the test level of $\alpha=0.05$, there is no statistical difference between the results of the two methods, and the results of the two evaluation methods are consistent, as shown in Table 3.

Table 3 Paired t-test results of two pain assessment methods

Method	mean	N	Std.Deviation	Std.Error Mean	t	P
FPS-R	3.68	41	1.724	0.269	1.871	0.77
POFM	3.44	41	1.534	0.240	1.817	0.77

Discussion. The different types of pain assessment scales currently used in clinical practice in China are translated into Chinese from English versions of scales used in European and American countries. However, the differences between Chinese and Western cultures, languages, habits, and medical systems, cause errors in the translation process, which directly affect the accuracy of pain assessment results and bring misjudgment to effective treatment (Wan Li, 2019 Li Jun, Feng Yi, 2011).

The pain measurement methods developed using samples may only come from one country, and / or from different cultures (for example, Western or Eastern); however, people from different cultural backgrounds speaking different languages may experience different forms of pain, and have different experiences and descriptions of pain^[8]. This possible variability is of great significance for evaluating the effectiveness of pain quality measures, which are directly translated into different languages without cultural adaptation. The measurement method developed using patient samples from one country or cultural background, directly translated into one's own language for use, may not necessarily be suitable for evaluating pain quality in other countries or cultures (Rahel Rogger, 2023).

One disadvantage of FPS-R is that subjects (patients) need to carefully observe and identify cartoon expressions before evaluation (Serife O, Aysenur P. et al., 2020). Because patients' pain levels are measured through selection of an image of a facial expression, the evaluation results are affected by cultural and other interference factors (Min Yoo, 2019). It is worth mentioning that the score of FPS-R is a categorical variable (an integer score of 0 to 10), and there are eleven levels of evaluation, so it can be statistically tested by parameters.

Chinese people of all ages are very familiar with POFM expressions, so find them easy to identify^[14]. This study selected six POFM expressions to assess patients' levels of pain. The six expressions correspond to pleasure, discomfort, slight pain, pain, severe pain, and unbearable pain respectively, and are consistent with the pain score scale 0, 2, 4, 6, 8, 10. When using POFM to evaluate pain, the patient selects the POFM expression which best corresponds to the degree of pain they are experiencing, and the assessor records the selection in terms of the scale value of the pain score.

In this study, the main forms of patient pain were limb pain caused by fracture, and shoulder pain caused by stroke hemiplegia. In some patients, pain was caused by hand trauma, cervical spondylosis, and localised pain by lumbar disc herniation.

The research results indicate that for Chinese patients, POFM is a sufficiently valid tool for pain measurement, with significant correlations with another commonly-used internationally-recognized pain scale, Wong-Baker FPS-R.

There were a few limitations to the study. Firstly, patients' preferences of the pain assessment scales were not evaluated. Also, the utility of POFM was only tested in a specific pain model, such as limb pain caused by fractures, and shoulder pain caused by stroke hemiplegia; therefore, the results cannot be generalized for all pain conditions within a given population. Before the formal use of various functional pain assessment scales is adopted in clinical practice, more in-depth research into the reliability, validity, and sensitivity of the scales needs to be carried out (Jan L, 2022).

Conclusions. We provided additional support for the effectiveness of the POFM scale in assessing the severity of pain in rehabilitation patients. Our conclusion is that the POFM scale is a suitable tool, even for patients with low educational levels, as it is easy to use, easy to understand, and does not require reading or writing skills, making it particularly suitable for Chinese patients. The first author of this article declares that there is no potential conflict of interest. The research funding comes from the author's own source.

REFERENCES

Chen Jun, Wang Jiang Lin. The international association for the study of pain (IASP) revision and systematic classification of chronic pain for WHO ICD-11. Chinese Journal of Pain Medicine. 2019; 25 (5):323-330. doi: 10.3969/j.issn.1006-9852 2019.05.002.

Michael P Rogers, Paul C Kuo. Pain as the Fifth Vital Sign. Journal of the American College of Surgeons. 2020; 231(5):601-602. doi: 10.1016/i. jamcollsurg. 2020 07508.

Wan Li, Zhao Qing, Chen Jun, et al, Expert consensus on the application of pain evaluation questionnaires in China (2020). Chinese Journal of Painology. 2020; 6(3):177-187. doi: 10.3760/cma.jcn101379-20190915-00075.

Davis MP, Walsh D. Cancer pain: how to measure the fifth vital sign. Clev Clin J Med. 2004;71(8):625-632. doi: 10.3949/ccjm.71.8.625.

Hicks CL, von Baeyer CL, Spafford PA, et al. The faces pain scale revised: toward a common metric in pediatric pain measurement. Pain. 2001;93(2):173-183. doi: 10.1016 / S0304-3959(01) 00314-1.

Thong IS, Jensen MP, Miró J, et al. The validity of pain intensity measures: what do the NRS, VAS, VRS, and FPS-R measure ? Scand J Pain. 2018; 18(1): 99-107. doi: 10.1515 / sjpain-2018- 0012.

Wan Li. Milestones in the compiling of pain related assessment scale of Chinese edition. Pain Clinic Journal. 2019;15(4):241. doi:10.3760/cma.j. issn.1672-9633.2019; 04.001.

Saurab Sharma, Anupa Pathak, Mark P Jensen. Words that describe chronic musculoskeletal pain: implications for assessing pain quality across cultures. Journal of Pain Research. 2016;9:1057-1066. doi:10.2147/jpr.s119212.

Ming Yan1, Rui Xiong, Yinghua Shen, et al. Intelligent generation of Peking opera facial masks with deep learning frameworks. Heritage Science. 2023;11 (20) 1-14. doi:10.1186/s40494-023-00865-z.

Fachao Zhang, Xiaoman Liang, Yaqi Sun, et al. POFMakeup: A style transfer method for Peking Opera makeup. Computers and Electrical Engineering. 2022;104 (PA):108459. doi: 10.1016/j.compeleceng.2022.108459.

Li Jun, Feng Yi, Han Ji Sheng, et al. Linguistic adaptation into Chinese and multicentered validation of the three questionnaires for the screening of neuropathic pain. Chinese Journal of Pain Medicine. 2011;17(9): 549-553. doi:10.3969/j.issn.1006-9852.2011.09.011.

Rahel Rogger, Corina Bello, Carolina S. Romero. Cultural Framing and the Impact On Acute Pain and Pain Services. Current Pain and Headache Reports. 2023;27:429-430. doi:10.1007/s11916-023-01125-2.

Min Yoo, Bao Ping, et al. Cross-cultural adaptation of the Neck Pain and Disability Scale: a methodological systematic review. The SPINE JOURNAL. 2019;19(6):1057-1066. doi: 10.1016/j.spinee.2019.01.007.

Fachao Zhang , Xiaoman Liang , et al. POFMakeup: A style transfer method for Peking Opera makeup. Computers and Electrical Engineering 2022; 104: 108459. doi: 10.1016/j.compeleceng.2022.108459.

Jan L, Thomas G, Jeffrey W. Validity and reliability of the Selective Control of the Upper Extremity Scale in children with upper motor neuron lesions. Disability and Rehabilitation. 2022; 44 (14): 3694-3700. doi: 10.1080/09638288.2021.1881171.

Serife O, Aysenur P. et al. Simulation of three intraoral radiographic techniques in pediatric dental patients: subjective comfort assessment using the VAS and Wong-Baker FACES Pain Rating Scale. BMC Oral Health. 2020; 20 (33):1-7. doi: 10.1186/s12903-020-1011-2. Serife O, Aysenur P. et al., 2020

<https://wongbakerfaces.org/us/wong-baker-faces-history/>.

Comprehensive evaluation of the functional state in senior schoolchildren with varying levels of daily motor activity / A. Martusevich, I. Bocharin, A. Eshiev [et al.] // Journal of Physical Education and Sport. – 2024. – Vol. 24, No. 11. – P. 1919-1926. – DOI 10.7752/jpes.2024.11288.

Improving the physical health of female students using boxing specialization in physical education / M. Kudryavtsev, V. Kovalev, A. Osipov [et al.] // Journal of Physical Education and Sport. – 2023. – Vol. 23, No. 7. – P. 1576-1582. – DOI 10.7752/jpes.2023.07193.

Results of using a plant adaptogen to improve the functional state of athletes / I. Bocharin, A. Eshiev, M. Guryanov [et al.] // Journal of Physical Education and Sport. – 2024. – Vol. 24, No. 11. – P. 1884-1891. – DOI 10.7752/jpes.2024.11284.

Correcting the physical health in 10–12-year-old schoolchildren through a creative approach to health-focused physical education / E. Romanova, H. Alshuwaili, G. Zhutanova [et al.] // Journal of Physical Education and Sport. – 2025. – Vol. 25, No. 2. – P. 284-291. – DOI 10.7752/jpes.2025.02032.

Engaging in sports as a method to enhance the stress resilience of a student's body / N. Gubareva, E. Romanova, A. Vorozheikin [et al.] // Journal of Physical Education and Sport. – 2024. – Vol. 24, No. 2. – P. 360-367. – DOI 10.7752/jpes.2024.02043.

Enhancing physical and functional fitness through the integration of health fitness elements in young gymnasts / E. V. Romanova, A. V. Vorozheikin, O. V. Bayankin [et al.] // Journal of Physical Education and Sport. – 2023. – Vol. 23, No. 10. – P. 2685-2691. – DOI 10.7752/jpes.2023.10307.

Personalization of physical education based on the study of body composition and body vegetative support in female students / M. Guryanov, I. Bocharin, E. Kalyuzhny [et al.] // Journal of Physical Education and Sport. – 2024. – Vol. 24, No. 2. – P. 353-359. – DOI 10.7752/jpes.2024.02042.

Physical fitness and life quality of female students with different classes of diseases / M. Kolokoltsev, E. Romanova, T. Martirosova [et al.] // Journal of Physical Education and Sport. – 2023. – Vol. 23, No. 1. – P. 128-133. – DOI 10.7752/jpes.2023.01016.

Bocharin, I., Eshiev, A., Guryanov, M., et al. Results of using a plant adaptogen to improve the functional state of athletes. Journal of Physical Education and Sport. 2024; 24(11): 1884-1891. doi: 10.7752/jpes.2024.11284.

Chen, J., Wang, J.L. The international association for the study of pain (IASP) revision and systematic classification of chronic pain for WHO ICD-11. Chinese Journal of Pain Medicine. 2019; 25(5): 323-330. doi: 10.3969/j.issn.1006-9852.2019.05.002.

Davis, M.P., Walsh, D. Cancer pain: how to measure the fifth vital sign. Cleveland Clinic Journal of Medicine. 2004; 71(8): 625-632. doi: 10.3949/ccjm.71.8.625.

Gubareva, N., Romanova, E., Vorozheikin, A., et al. Engaging in sports as a method to enhance the stress resilience of a student's body. Journal of Physical Education and Sport. 2024; 24(2): 360-367. doi: 10.7752/jpes.2024.02043.

Guryanov, M., Bocharin, I., Kalyuzhny, E., et al. Personalization of physical education based on the study of body composition and body vegetative support in female students. Journal of Physical Education and Sport. 2024; 24(2): 353-359. doi: 10.7752/jpes.2024.02042.

Hicks, C.L., von Baeyer, C.L., Spafford, P.A., et al. The faces pain scale revised: toward a common metric in pediatric pain measurement. *Pain*. 2001; 93(2): 173-183. doi: 10.1016/S0304-3959(01)00314-1.

Jan, L., Thomas, G., Jeffrey, W. Validity and reliability of the Selective Control of the Upper Extremity Scale in children with upper motor neuron lesions. *Disability and Rehabilitation*. 2022; 44(14): 3694-3700. doi: 10.1080/09638288.2021.1881171.

Kolokoltsev, M., Romanova, E., Martirosova, T., et al. Physical fitness and life quality of female students with different classes of diseases. *Journal of Physical Education and Sport*. 2023; 23(1): 128-133. doi: 10.7752/jpes.2023.01016.

Kudryavtsev, M., Kovalev, V., Osipov, A., et al. Improving the physical health of female students using boxing specialization in physical education. *Journal of Physical Education and Sport*. 2023; 23(7): 1576-1582. doi: 10.7752/jpes.2023.07193.

Li, J., Feng, Y., Han, J.S., et al. Linguistic adaptation into Chinese and multicentered validation of the three questionnaires for the screening of neuropathic pain. *Chinese Journal of Pain Medicine*. 2011; 17(9): 549-553. doi: 10.3969/j.issn.1006-9852.2011.09.011.

Martusevich, A., Bocharin, I., Eshiev, A., et al. Comprehensive evaluation of the functional state in senior schoolchildren with varying levels of daily motor activity. *Journal of Physical Education and Sport*. 2024; 24(11): 1919-1926. doi: 10.7752/jpes.2024.11288.

Rogers, M.P., Kuo, P.C. Pain as the Fifth Vital Sign. *Journal of the American College of Surgeons*. 2020; 231(5): 601-602. doi: 10.1016/j.jamcollsurg.2020.07.508.

Rogger, R., Bello, C., Romero, C.S. Cultural Framing and the Impact On Acute Pain and Pain Services. *Current Pain and Headache Reports*. 2023; 27: 429-430. doi: 10.1007/s11916-023-01125-2.

Romanova, E., Alshuwaili, H., Zhutanova, G., et al. Correcting the physical health in 10–12-year-old schoolchildren through a creative approach to health-focused physical education. *Journal of Physical Education and Sport*. 2025; 25(2): 284-291. doi: 10.7752/jpes.2025.02032.

Serife, O., Aysenur, P., et al. Simulation of three intraoral radiographic techniques in pediatric dental patients: subjective comfort assessment using the VAS and Wong-Baker FACES Pain Rating Scale. *BMC Oral Health*. 2020; 20(33): 1-7. doi: 10.1186/s12903-020-1011-2.

Sharma, S., Pathak, A., Jensen, M.P. Words that describe chronic musculoskeletal pain: implications for assessing pain quality across cultures. *Journal of Pain Research*. 2016; 9: 1057-1066. doi: 10.2147/jpr.s119212.

Thong, I.S., Jensen, M.P., Miró, J., et al. The validity of pain intensity measures: what do the NRS, VAS, VRS, and FPS - R measure? *Scandinavian Journal of Pain*. 2018; 18(1): 99 - 107. doi: 10.1515/sjpain-2018-0012.

Wan, L., Zhao, Q., Chen, J., et al. Expert consensus on the application of pain evaluation questionnaires in China (2020). *Chinese Journal of Painology*. 2020; 6(3): 177-187. doi: 10.3760/cma.j.cn101379-20190915-00075.

ISSN 2414-0244

Научно-периодический журнал «Здоровье человека, теория и методика физической культуры и спорта». - 2026. - 41 (1)

Раздел 2. МЕДИКО-БИОЛОГИЧЕСКИЕ ПРОБЛЕМЫ ЗДОРОВЬЯ ЧЕЛОВЕКА

DOI: [https://doi.org/10.14258/zosh\(2026\)1.04](https://doi.org/10.14258/zosh(2026)1.04)

Wan, L. Milestones in the compiling of pain related assessment scale of Chinese edition. *Pain Clinic Journal*. 2019; 15(4): 241. doi: 10.3760/cma.j.issn.1672-9633.2019.04.001.

Wong-Baker FACES Pain Rating Scale. History. Available from: <https://wongbakerfaces.org/us/wong-baker-faces-history/>

Yan, M., Xiong, R., Shen, Y., et al. Intelligent generation of Peking opera facial masks with deep learning frameworks. *Heritage Science*. 2023; 11(20): 1-14. doi: 10.1186/s40494-023-00865-z.

Yoo, M., Bao, P., et al. Cross-cultural adaptation of the Neck Pain and Disability Scale: a methodological systematic review. *The Spine Journal*. 2019; 19(6): 1057-1066. doi: 10.1016/j.spinee.2019.01.007.

Zhang, F., Liang, X., Sun, Y., et al. POFMakeup: A style transfer method for Peking Opera makeup. *Computers and Electrical Engineering*. 2022; 104(PA): 108459. doi: 10.1016/j.compeleceng.2022.108459.